



HYPOGLYCAEMIC EFFECT OF VRIDDHADARU IN ALLOXAN INDUCED DIABETIC RABBITS

Vivek P¹, Jayakumari D², Jayasree P³

¹Assistant professor, VPSV Ayurveda College, Kottakkal, Kerala, email – drvivekp@gmail.com

²Professor, Maria Ayurveda Medical College, Marthandam, Tamilnadu,

³Professor, Govt Ayurveda College, Thiruvananthapuram, Kerala Email - drjayasree11@yahoo.com

ABSTRACT

Diabetes Mellitus is the most important non communicable disease, playing a notorious role in the devastating phase of public health. Ayurveda, the life science provides astonishing results in such life style disorders. Vriddhadaru an excellent Ayurvedic drug and *Argyreia nervosa* has been traditionally used as Vriddhadaru, effective in diabetic management with promising results. But, the anti-diabetic effect of the drug has not been scientifically appreciated. This study ascertains the hypoglycaemic effect of the drug in a pre-clinical model. In this study, alloxan induced 18 diabetic albino rabbits were divided into 3 groups with 6 rabbits in each group. On the day of experiment, after assessing the fasting blood sugar levels in all 18 rabbits, root powder of the test drug as CMC suspension in the dose of 0.56 g/kg body weight and the standard drug, metformin; in the dose of 0.024 mg/kg body weight were administered to the first group and second group respectively. The third group was kept as control and administered with distilled water alone. Later, the blood glucose values were taken at 1st, 3rd and 5th hour and the obtained values were compared within the group and between the groups. The blood glucose levels were compared using paired t test for within the group analysis and by student t test for between the group analysis. Within the group comparison of blood sugar level in group 1 showed significant decrease in the level at 1st hour and 5th hour from the corresponding fasting blood sugar levels. In group 2, significant decrease of blood sugar level was seen between fasting blood sugar level and 1st hour value and also in fasting blood sugar value and 5th hour value. Between group comparison of group 2 showed significant difference in the reducing the blood sugar level at different time intervals. Ayurvedic literature proposes Pramehaghna property to the drug Vriddhadaru which is substantiated through this pre-clinical trial. But the effect is comparatively less with that of the standard drug. This may be due to the absence of strong antioxidant and antidiabetic chemical constituents in this plant. From the above study, it is inferred that *Argyreia nervosa* (Burm.f.) Boj. is less effective in reducing blood sugar levels in Alloxan induced diabetic rabbits against the standard drug Metformin.

Key words: *Alloxan monohydrate, Argyreia nervosa*

INTRODUCTION

Life style disorders and non-communicable diseases causing life distress, melancholy and untimely death now a days. Among all non-communicable diseases, metabolic syndrome

stands apart and is of quite a concern. This condition mainly includes dyslipidemia and diabetes mellitus type II. The latter is characterised by chronic hyperglycaemia with disturbances of carbohydrate, fat and protein metabolism resulting from defects in insulin secretion, insulin action, or both. It occurs when the pancreas does not produce enough insulin, or when the body cannot effectively use the insulin it produces¹.

Diabetes is in the top 10, and perhaps the top 5, of the most significant diseases in the developed world, and is gaining significance elsewhere. In 2006, according to the WHO, at least 171 million people worldwide suffer from diabetes. Its incidence is increasing rapidly, and it is estimated that by the year 2030, this number will double². The effects of diabetes mellitus include long-term damage, dysfunction and failure of various organs. In its most severe forms, ketoacidosis or a non- ketotic hyperosmolar state may develop and lead to stupor, coma and, in absence of effective treatment, death. Even though a large number of drugs are available for this condition, none ensure complete cure, especially in preventing complications³.

Many traditionally using Ayurvedic drugs were possessed pramehaghna property still many of them are having no scientifically validated data. Vriddhadaru is one if such drug mentioned firstly in Rasayana prakarana of Ashtanga Samgraha⁴. Even though many controversies are there about the identification of the genuine source plant, *Argyreia nervosa* (Burm. f.) Boj. of Convolvulaceae family is considered as the mostly used plant source as Vriddhadaru⁵. *Argyreia nervosa* is a climbing shrub with woody tomentose stem, commonly known as elephant creeper in English. It is widely distributed in tropical regions of the world. In India it is seen upto an altitude of 900 m⁶. It is generally found growing in slightly moist localities like river banks, edges of lakes etc. The roots are varying in size as well as in thickness⁷.

As per Ayurvedic classical texts, this plant is useful in Prameha⁸, a condition in which a drug Pramehaghna with Rasayana property is very much essential. This aspect of Vriddhadaru has not been established till date. Moreover, search for new effective drugs are very much essential now a days as genuine Ayurvedic drugs in this picture are facing radical scarcity.

There are many studies are there about the analgesic^{9,10}, anti-inflammatory^{11,12}, antipyretic¹³, antiviral¹⁴, anticulvulant¹⁵, nootropic^{16,17}, anthelmintic¹⁸, aphrodisiac^{19,20,21}, antidiarrheal²², antiulcer²³ and antimicrobial action^{24,25,26}. But only a few are there about the hypoglycaemic activity. Hence in this study, it was decided to take *Argyreia nervosa* as a source plant for Vriddhadaruka and to document the hypoglycaemic activity in alloxan induces diabetic rabbits against the standard drug metformin.

MATERIALS AND METHODS

a) Selection of animals

25 healthy albino rabbits of either sexes weighing 1500-2000gm were collected from the animal house of Agadatantra department, Government Ayurveda College, Thiruvananthapuram.

b) Preparation of Alloxan solution

Alloxan monohydrate was obtained from Laboratory supplies Thiruvananthapuram, batch no: 43256. Using normal saline 0.5%, the solution was prepared.

c) **Standard drug - Metformin** - A stock solution was prepared with 0.5 mg of metformin in 50 ml distilled water so that 1 ml contained 0.02 ml of metformin²⁷

d) Preparation of test drugs

Form of preparation- Powder

Route of administration- Oral

Drug- Root of *Argyreia nervosa* (Burm. f.) Boj.

A Carboxymethylcellulose stock solution of strength 0.01g% of the test drug was prepared in distilled water.

e) Dose

The dose of the test drug was calculated from Sarngdhara samhitha; the dose being the common dose for choorna²⁸. Effective dose for rabbit was calculated by using the table constructed by Paget G.E. and Barnes T.M. in the evaluation of drug activities. Based on this; corresponding doses were calculated.

f) Setting – Animal house working under the Drug Standardization Unit, Govt. Ayurveda College, Trivandrum, Kerala

g) Procedure²⁹ - 25 albino rabbits of either sex (weighing 1500-2000gm) were used in the present study. The animals were kept under observation for one week. Normal fasting blood sugar values were noted before administration of alloxan. All rabbits were fasted for 18 hours and the samples were collected from marginal ear vein. Blood sugar was calculated by glucose peroxidase (GOD-POD) method. After that, normal rabbit feed was given. Alloxan monohydrate (150 mg/kg body weight) dissolved in normal saline was injected via intra peritoneal route in 18 h previously fasted animal to induce diabetes. After one hour of alloxan administration, the animals were fed with standard pellets and water *ad libitum*. After 72 hours, the blood glucose levels were estimated, applying the glucose oxidase method and rabbits having blood glucose level more than double were selected for the study³⁰.

18 healthy diabetic rabbits were taken for the study. They were grouped into three, with each group containing 6 animals which were caged separately. Animals of each group were marked for individual identification. First group (G1) was treated with test drug i.e. 0.56 g/kg body weight of root powder of *Argyreia nervosa* (Burm. f.) Boj., second (G2) group was administered with standard drug metformin at the dose of 0.024 mg/kg and the third group (G3) was kept as control and treated with distilled water alone.

On the day of the experimental study, fasting blood samples were taken from all the rabbits and then; the drugs were administered orally using feeding cannula. After 1 hour blood samples were collected and all the animals were fed. Blood samples for glucose assessment were again collected from all animals at 3rd and 5th hours after the drug administration. The values were subjected to statistical analysis.

OBSERVATIONS AND RESULTS

The blood glucose levels were compared using paired t test for within the group analysis and by student t test for between the group analysis using the programme WinPepi version 11. While comparing the blood sugar values in G1, using paired t, the following data were obtained

Table No. 1

	Mean	SD	Mean diff	p - Value
FBS and 1 st hour	232.0 195.7	29.4 12.7	36.2	< 0.01
FBS and 3 rd hour	232.0 217.9	29.4 16.6	14.0	> 0.05
FBS and 5 th hour	232.0 186.7	29.4 19.1	45.2	< 0.01
1 st hour and 3 rd hour	195.7 217.9	12.7 16.6	22.2	< 0.05
1 st hour and 5 th hour	195.7 186.7	12.7 19.1	9.0	> 0.05
3 rd hour and 5 th hour	217.9 186.7	16.6 19.1	31.2	< 0.01

The comparison of the blood sugar level at different time of G2, using paired t test is given below.

Table No. 2

	Mean	SD	Mean diff	p - Value
FBS and 1 st hour	246.6 161.4	34.1 37.9	85.2	< 0.01
FBS and 3 rd hour	246.6 212.4	34.1 31.0	34.1	> 0.05
FBS and 5 th hour	246.6 150.2	34.1 34.3	96.4	< 0.01
1 st hour and 3 rd hour	161.4 212.4	37.9 31.0	51.0	< 0.01
1 st hour and 5 th hour	161.4 150.2	37.9 34.3	11.2	> 0.05
3 rd hour and 5 th hour	212.4 150.2	31.0 34.3	62.3	< 0.01

The comparison of the blood sugar level at different time of G3, using paired t test is given below.

Table No. 3

	Mean	SD	Mean diff	p - Value
FBS and 1 st hour	252.6 233.1	26.0 21.7	19.5	< 0.05
FBS and 3 rd hour	252.6 291.2	26.0 21.0	38.6	< 0.05
FBS and 5 th hour	252.6 273.4	26.0 18.5	20.8	> 0.05
1 st hour and 3 rd hour	233.1 291.2	21.7 21.0	58.1	< 0.01
1 st hour and 5 th hour	233.1 273.4	21.7 18.5	40.3	< 0.05
3 rd hour and 5 th hour	291.2 273.4	21.0 18.5	17.8	< 0.01

Comparison of average changes in blood sugar level between group 1 and group 2 was done using unpaired t test and the results were given in the table number 4

Table No. 4

Duration	Group 1		Group 2		p value
	Mean	SD	Mean	SD	
FBS & 1 st hour	14.91	7.42	34.45	12.83	< 0.01
FBS & 3 rd hour	5.13	10.23	13.07	13.54	> 0.05
FBS & 5 th hour	19.01	7.12	38.38	14.98	< 0.05
1 st hour & 3 rd hour	11.52	7.57	33.86	12.84	< 0.01
1 st hour & 5 th hour	4.51	8.30	5.92	16.09	> 0.05
3 rd hour & 5 th hour	14.39	4.67	29.45	11.98	< 0.05

Within the group comparison of blood sugar level in group 1 showed significant decrease in the level at 1st hour ($p < 0.01$) and 5th hour ($p < 0.01$) from the corresponding fasting blood sugar levels. No significant difference was there between fasting blood sugar level and 3rd hour value ($p > 0.05$). Significant increase in the blood sugar level was noted from 1st hour to 3rd hour ($p < 0.05$) then gradually decreased, but not statistically significantly ($p > 0.05$) while comparing it with 1st hour value. A significant decrease at $p < 0.01$ was observed between 3rd hour and 5th hour blood glucose level.. Hence, initially there was a significant decrease in the blood sugar at 1st hour. Then in 3rd hour there was a significant increase of value than that of 1st hour followed by a gradual decrease in 5th hour.

In group 2, significant decrease of blood sugar level was seen between fasting blood sugar level & 1st hour value ($p < 0.01$) and fasting blood sugar value & 5th hour value ($p < 0.01$). No significant change was observed between fasting blood sugar level and 3rd hour value ($p > 0.05$). Significant increase in the blood sugar level was noted between 1st and 3rd hour value ($p < 0.01$) but no difference was observed between 1st and 5th hour value ($p > 0.05$). But 3rd and 5th values are different significantly ($p < 0.01$).

In group 3, the blood sugar levels increased significantly ($p < 0.05, 0.01$) in most of the cases and in few cases there was no statistical difference ($p > 0.05$)

Between group comparison of group 1 and group 2 showed significant difference in the change of the blood sugar level at different time intervals. Mean values showed that standard drug is more effective than the test drug.

DISCUSSION

Statistical evaluation shows that the blood glucose level in group 1, treated with *Argyreia nervosa* shows a gradual decrease but less effective than that of standard drug metformin.

Previous studies show that *Argyreia nervosa* contains 1-tricontanol, epifriedelinol acetate, epifriedelinol and β -sitosterol²⁵ and the hexane extract of the root yielded tetradecanyl palmitate, 5, 8-oxidotetracosan-10-one²⁶ and two novel aryl esters characterized as stigmasteryl phydroxycinnamate and hexadecanyl p-hydroxycinnmate along with scopoletin²⁶. Even though some of these were told to be having antioxidant, a prominent hypoglycaemic activity has not proven for any of this constituent. This may be the reason for the weak hypoglycaemic activity than the standard drug.

CONCLUSION

From the above study, it is inferred that *Argyreia nervosa* (Burm.f) Boj. is effective in reducing the blood glucose level in reducing blood sugar levels in Alloxan induced diabetic rabbits but less effective while comparing with that of the standard drug Metformin. Effectiveness of the plant and advantages if any over the existing therapy can only be assessed by a long term study in more than one type of rodents. Clinical efficacy in human subjects should also need to be proved for establishing it as an effective single drug therapy in Diabetes mellitus.

REFERENCES

1. WHO, Diabetes, Available from Netlibrary http://www.who.int/topics/diabetes_mellitus/en/
2. WHO, Global status report on noncommunicable diseases 2014, Available from Netlibrary <http://www.who.int/nmh/publications/ncd-status-report-2014/en/>
3. Fauci et al., 2007, Harrison's principles of internal medicine, 17th edition, Mc Grow Hill Publication, pp. 2275
4. Vagbhata, Astanga samgraha, Uttarasthaana, translated by Srikantha Murthy K R, Choukhambha orientalia, Varanasi, Vol 3, 49/342, pp: 510
5. Aiyer KN, Kolammal M. Pharmacognosy of Ayurvedic Drugs Kerala, 1(8), Department of Pharmacognosy, University of Kerala, Trivandrum, 1964, pp. 61-65.
6. Anonymous, Flora of Orissa. Orissa forest development co. Ltd; Bhubaneswar, Orissa, 1995.
7. Gamble JS. Flora of Madras. Botanical survey of, India, Calcutta, volume II, 1956, Reprint, 556.
8. Vagbhata, Astanga samgraha, Uttarasthaana, translated by Srikantha Murthy K R, Choukhambha orientalia, Varanasi, Vol 3, 49/344, pp: 510
9. Bachhav RS, Gulecha VS, Upasani CD. Analgesic and inflammatory activity of *Argyreia speciosa* root. Ind J Pharmacol. 2009, 41, 158-61
10. Varsh J, Galani, Bharatkumar P. Analgesic and anti-inflammatory activity of *Argyreia nervosa* and *Sphearanthus indicus* in the experimental animals Global J Pharmacol. 2010, 4(3), 136-141.
11. Srivastava MC, Kant V, Tewari JP. Antiinflammatory activity of roots of *Argyreia speciosa* (Sumandrashokha). Mediscope. 1972, 15, 219-222.
12. Bachhav RS, Gulecha VS, Upasani CD. Analgesic and inflammatory activity of *Argyreia speciosa* root. Ind J Pharmacol. 2009, 41, 158-61.
13. Kamal J, Sunil T, Narender T. Antipyretic activity of whole aerial part from *Argyreia nervosa*. Int J Pharma Pharmaceut Sci. 2012, 4 (4), 1-2.
14. Babbar OP, Joshi MN, Madan AR. Evaluation of plants for antiviral activity. Ind J Med Res. 1982, 76, 54-65.
15. Vyawahare NS, Bodhankar SL. Anticonvulsant activity of *Argyreia speciosa* in mice. Ind J Pharm Sci. 2009, 71, 131-134.
16. Joshi H, Kaur N, Chauhan J. Evaluation of Nootropic effect of *Argyreia speciosa* in mice. J Health Sci. 2007, 53, 382-388.
17. Vyawahare NS, Bodhankar SL. Effect of *Argyreia speciosa* extract on learning and memory paradigms in mice. Phcog Mag. 2009, 4, 43-48.
18. Parveen N, Khan NU, Singhal KC. Antifilarial activity of *Argyreia speciosa* against *Setaria cervi* in vitro. Phytother Res. 1990, 4, 162-164.

19. Subramoniam A, Madhavachandran V, Ravi K, Anju VS. Aphrodisiac property of the elephant creeper *Argyreia nervosa*. J Endocrinol Reprod. 2007, 11, 82- 85.
20. Jayatilak PG, Sheth AR, Mugatwala PP, Pardani DS. Effect of an indigenous drug (Speman) on human accessory reproductive function. Ind J Surg. 1976, 38, 12-15.
21. Bhargava NC, Singh OP. Fortege and indigenous drug in common sexual disorders in males. Mediscope. 1978, 21, 140-144.
22. Rao CV, Ojha SK, Reddy GD, Rawat AK, Rao GM, Pushpangadan P. Antidiarrhoeal activity of *Argyreia speciosa* flower: An ethanopharmacological study. Acta Pharm Turcica. 2004, 46, 149-159.
23. Khan A, Roshan S, Tazneem B, Ali S. Anti-ulcer effect of *Argyreia speciosa* ethanolic root extract in rats. Pharmacologyonline. 2010, 2, 389-394.
24. Ashish JM, Khadabadi SS, Farooqui IA, Ghorpade DS. Studies on antimicrobial activity of *Clerodendrum infortunatum*, *Argyreia nervosa* and *Vitex negundo*: A comparison. Der Pharma Lett. 2010, 2 (1), 102-105.
25. Basha SK, Sudarshanam G, Evaluation of antibacterial activity of some Medicinal plants used by Sugali tribe of Yerramalais forest of Andhra Pradesh. J Pharma Res. 2011, 4(7), 2067-2068.
26. Somashekhar K, Ajay M, Pallavi R, Deepali SG. Invitro antifungal activity of ethanol fraction of *Argyreia nervosa* (Burm. f.) Boj. Leaves. Ind J Nat Prod Res. 2012, 3 (1), 48-54.
27. Gerhard Vogel H, Drug discover and evaluation pharmacological assays, Springer, second edition, 2002, pp:950
28. Sargadhara, Sarngadhara Samhitha, English translation by Srikantha Murthy K. R., Chaukhambha orientalia, Varanasi, sixth edition 2004, 6/1, pp 84.
29. Gerhard Vogel H, Drug discover and evaluation pharmacological assays, Springer, second edition, 2002, pp:950
30. Vijay S Patel, V Chitra, P Lakshmi Prasanna, V Krishnaraju, Hypoglycemic effect of aqueous extract of *Parthenium hysterophorus* L. in normal and alloxan induced diabetic rats Department of Pharmacology, SRM College of Pharmacy, Indian journal of Pharmacology, 2008 Volume : 40 Issue : 4 P : 183-185, Available from net libraray <http://www.ijp-online.com/article.asp?issn=0253-13;year=2008;volume=40;issue=4;spage=183;epage=185;aulast=Patel>.